




**Romania**  
**Team Selection Tests**  
Timisoara, Bucharest 2008



**Day 1 - 01 May 2008**

- 1] Let  $n$  be an integer,  $n \geq 2$ . Find all sets  $A$  with  $n$  integer elements such that the sum of any nonempty subset of  $A$  is not divisible by  $n + 1$ .
- 2] Let  $a_i, b_i$  be positive real numbers,  $i = 1, 2, \dots, n$ ,  $n \geq 2$ , such that  $a_i < b_i$ , for all  $i$ , and also

$$b_1 + b_2 + \dots + b_n < 1 + a_1 + \dots + a_n.$$

Prove that there exists a  $c \in \mathbb{R}$  such that for all  $i = 1, 2, \dots, n$ , and  $k \in \mathbb{Z}$  we have


$$(a_i + c + k)(b_i + c + k) > 0.$$

- 3] Let  $ABCDEF$  be a convex hexagon with all the sides of length 1. Prove that one of the radii of the circumcircles of triangles  $ACE$  or  $BDF$  is at least 1.
- 4] Prove that there exists a set  $S$  of  $n - 2$  points inside a convex polygon  $P$  with  $n$  sides, such that inside any triangle determined by 3 vertices of  $P$  contains exactly one point from  $S$  inside or on the boundaries.
- 5] Find the greatest common divisor of the numbers

$$2^{561} - 2, 3^{561} - 3, \dots, 561^{561} - 561.$$



**Romania**  
**Team Selection Tests**  
Timisoara, Bucharest 2008



**Day 2**

- 1] Let  $n \geq 3$  be an odd integer. Determine the maximum value of

$$\sqrt{|x_1 - x_2|} + \sqrt{|x_2 - x_3|} + \dots + \sqrt{|x_{n-1} - x_n|} + \sqrt{|x_n - x_1|},$$

where  $x_i$  are positive real numbers from the interval  $[0, 1]$ .


- 2] Are there any sequences of positive integers  $1 \leq a_1 < a_2 < a_3 < \dots$  such that for each integer  $n$ , the set  $\{a_k + n \mid k = 1, 2, 3, \dots\}$  contains finitely many prime numbers?
- 3] Show that each convex pentagon has a vertex from which the distance to the opposite side of the pentagon is strictly less than the sum of the distances from the two adjacent vertices to the same side.

*Note.* If the pentagon is labeled  $ABCDE$ , the adjacent vertices of  $A$  are  $B$  and  $E$ , the ones of  $B$  are  $A$  and  $C$  etc.

- 4] Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges such that each edge is contained in at least one triangle. Find the minimum value of  $m$ .



**Romania**  
**Team Selection Tests**  
Timisoara, Bucharest 2008



**Day 3**

- 1] Let  $ABC$  be a triangle with  $\angle BAC < \angle ACB$ . Let  $D, E$  be points on the sides  $AC$  and  $AB$ , such that the angles  $ACB$  and  $BED$  are congruent. If  $F$  lies in the interior of the quadrilateral  $BCDE$  such that the circumcircle of triangle  $BCF$  is tangent to the circumcircle of  $DEF$  and the circumcircle of  $BEF$  is tangent to the circumcircle of  $CDF$ , prove that the points  $A, C, E, F$  are concyclic.

*Author: Cosmin Pohoata*


- 2] Let  $ABC$  be an acute triangle with orthocenter  $H$  and let  $X$  be an arbitrary point in its plane. The circle with diameter  $HX$  intersects the lines  $AH$  and  $AX$  at  $A_1$  and  $A_2$ , respectively. Similarly, define  $B_1, B_2, C_1, C_2$ . Prove that the lines  $A_1A_2, B_1B_2, C_1C_2$  are concurrent.

*Remark.* The triangle obviously doesn't need to be acute.

- 3] Let  $m, n \geq 3$  be positive odd integers. Prove that  $2^m - 1$  doesn't divide  $3^n - 1$ .
- 4] Let  $n$  be a nonzero positive integer. A set of persons is called a  $n$ -balanced set if in any subset of 3 persons there exists at least two which know each other and in each subset of  $n$  persons there are two which don't know each other. Prove that a  $n$ -balanced set has at most  $(n - 1)(n + 2)/2$  persons.



**Romania**  
**Team Selection Tests**  
Timisoara, Bucharest 2008




**Day 4 - 12 June 2008**

- 1] Let  $ABCD$  be a convex quadrilateral and let  $O \in AC \cap BD$ ,  $P \in AB \cap CD$ ,  $Q \in BC \cap DA$ . If  $R$  is the orthogonal projection of  $O$  on the line  $PQ$  prove that the orthogonal projections of  $R$  on the sidelines of  $ABCD$  are concyclic.
- 2] Let  $m, n \geq 1$  be two coprime integers and let also  $s$  an arbitrary integer. Determine the number of subsets  $A$  of  $\{1, 2, \dots, m + n - 1\}$  such that  $|A| = m$  and  $\sum_{x \in A} x \equiv s \pmod{n}$ .
- 3] Let  $n \geq 3$  be a positive integer and let  $m \geq 2^{n-1} + 1$ . Prove that for each family of nonzero distinct subsets  $(A_j)_{j \in \overline{1, m}}$  of  $\{1, 2, \dots, n\}$  there exist  $i, j, k$  such that  $A_i \cup A_j = A_k$ .



**Romania**  
**Team Selection Tests**  
Timisoara, Bucharest 2008



**Day 5 - 13 June 2008**

- 1] Let  $n$  be a nonzero positive integer. Find  $n$  such that there exists a permutation  $\sigma \in S_n$  such that

$$|\{\sigma(k) - k : k \in \overline{1, n}\}| = n.$$

- 2] Let  $ABC$  be a triangle and let  $\mathcal{M}_a, \mathcal{M}_b, \mathcal{M}_c$  be the circles having as diameters the medians  $m_a, m_b, m_c$  of triangle  $ABC$ , respectively. If two of these three circles are tangent to the incircle of  $ABC$ , prove that the third is tangent as well.
- 3] Let  $\mathcal{P}$  be a square and let  $n$  be a nonzero positive integer for which we denote by  $f(n)$  the maximum number of elements of a partition of  $\mathcal{P}$  into rectangles such that each line which is parallel to some side of  $\mathcal{P}$  intersects at most  $n$  interiors (of rectangles). Prove that

$$3 \cdot 2^{n-1} - 2 \leq f(n) \leq 3^n - 2.$$